



NASOPHARYNGEAL CANCER IN THE WORLD: EPIDEMIOLOGY, INCIDENCE, MORTALITY AND RISK FACTORS

H. SALEHINIYA^{1,2}, M. MOHAMMADIAN³, A. MOHAMMADIAN-HAFSHEJANI⁴,
N. MAHDAVIFAR³

¹Zabol University of Medical Sciences, Zabol, Iran

²Department of Epidemiology and Biostatistics, School of Public Health,
Tehran University of Medical Sciences, Tehran, Iran

³Health Promotion Research Center, Department of Epidemiology and Biostatistics, School of Public Health,
Zahedan University of Medical Sciences, Zahedan, Iran

⁴Department of Epidemiology and Biostatistics, School of Public Health, Shahrekord University of Medical
Sciences, Shahrekord, Iran

Abstract – Objective: Nasopharyngeal cancer is one of the most common cancers in the head and neck areas. Men are two to three times more likely to develop it than women. The peak age of incidence is between 50 and 60 years. Informing about the occurrence, death and risk factors in prevention programs is very important. The aim of this study was to investigate the incidence, mortality, and risk factors for nasopharyngeal cancer in the world.

Materials and Methods: This review study was conducted on published English studies by January 2017 by searching in the databases of PubMed, Scopus and web of science. The search strategy included the key words “nasopharyngeal cancer”, “epidemiology”, “incidence”, “mortality”, “risk factor”, “world”. Studies related to incidence, mortality and risk factors for nasopharyngeal cancer were entered into the review.

Results: Nasopharyngeal cancer is generally more common in Southeast Asia. The five countries with the highest incidence of nasopharyngeal cancer in the world were China, Indonesia, Vietnam, India and Malaysia, respectively. The standard incidence of nasopharyngeal cancer in the world was 1.2 per 100,000 (in men 1.7 per 100,000; in women, 0.7 per 100,000). The five countries with the highest number of deaths were China, Indonesia, Vietnam, India and Malaysia, respectively. The standardized mortality rate for nasopharyngeal cancer in the world was 0.7 per 100,000 (in men 1.0 per 100,000; in women 0.4 per 100,000). The most important risk factors for nasopharyngeal cancer were Epstein-Barr virus (EBV), heredity, human leukocyte antigen (HLA) genes, salt-preserved fish consumption, and history of respiratory diseases.

Conclusions: The incidence of nasopharyngeal cancer is related to the geographic region. Training programs, early screening, good life promotion policies, timely treatment and diagnosis are good for reducing the burden of this cancer.

KEYWORDS: Nasopharyngeal Cancer, Incidence, Mortality, Risk Factor, World.

INTRODUCTION

One of the most common cancers in the head and neck is nasopharyngeal cancer^{1,2}. It is a malignant disease and its incidence is dependent on geographical and racial variations. Nasopharyngeal carcinoma has been reported in most parts of the world with age standardi-

zed incidence rate, regardless of gender, less than one per 100,000 per year³⁻⁵. This cancer has an unbalanced geographical distribution. 81% of the new cases occurred in Asia and 9% in Africa; the rest were reported elsewhere in the world. Southeast Asian countries account for 67% of the global burden of cancer⁶. In addition to geographic variation, some ethnic groups



may be at risk of nasopharyngeal cancer. For example: Bidayuh on the island of Borneo, Nagas in Northern India and Inuits in the Arctic, with an age-old standard of more than 16 per 100,000 per year in men⁷.

Men are two to three times more likely to develop it than women. The peak age of incidence is between 50 and 60 years⁸. Elderly people are at higher risk for relapse and also have a lower survival rate⁹. On the other hand, the highest mortality rate has been observed in people over the age of 85 years¹⁰. Diagnosis of nasopharyngeal cancer occurs at advanced stages. The prognosis for advanced nasopharyngeal cancer is very weak. However, nasopharyngeal cancers are potentially curable in the early stages. Identification in the early stages of screening may lead to improved results. Informing about the occurrence, death, and risk factors in prevention programs is very important. The aim of this study was to investigate the incidence, mortality, and risk factors for nasopharyngeal cancer in the world.

MATERIALS AND METHODS

This review study was conducted on published English studies by January 2017 by searching in the databases of PubMed, Scopus and web of science. The search strategy included the key words “nasopharyngeal cancer”, “epidemiology”, “incidence”, “mortality”, “risk factor”, “world”. Studies related to incidence, mortality and risk factors for nasopharyngeal cancer were entered into the review.

RESULTS

NUMBER AND INCIDENCE

In 2012, there were 86,691 cases of nasopharyngeal cancer worldwide (60,896 cases in men and 25,795 cases in women, Sex Ratio = 2.36). The standard incidence of nasopharyngeal cancer in the world was 1.2 per 100,000 (1.7 per 100,000 in men and 0.7 per 100,000 in women)¹¹. The five countries with the highest number of nasopharyngeal cancer cases inclu-

ded China with 42,100 cases¹², Indonesia with 13,084 cases, Vietnam with 4,931 cases, India with 3,947 and Malaysia with 2,030 cases, respectively¹³. In Singapore, nasopharyngeal cancer is the 8th common cancer in men, with the age-standardized rate of 9.5 per 100,000 per year¹⁴. In Indonesia, a relatively high incidence, at least 5.7 per 100,000 in males and 1.9 in women per 100,000 compared to the global incidence rate of 1.9 per 100,000 in men and 0.8 per 100,000 in women, has been reported¹⁵. It should be noted that the correct incidence of nasopharyngeal cancer in Indonesia is not clear due to incomplete cancer recording¹⁶. The lowest incidence of nasopharyngeal cancer has been reported significantly from the United States, America and Europe. Nasopharyngeal cancers in Europe and North America include less than 1% of all cases of cancer (17). This geographic distribution indicates the difference in the pathology of the nasopharyngeal cancer and its epidemiology in these areas^{17,18}. The high prevalence of nasopharyngeal cancer in these countries is due to exposure to all types of risk factors associated with this cancer and to lowering health budgets. Diagnosis in the advanced stage and the lack of access to treatment, metastatic state of the cancer is more observed in these areas^{19,20}.

NUMBER AND MORTALITY RATE

In the world in 2012, 50,831 deaths from nasopharynx occurred (35,756 deaths in men and 15075 deaths in women, Sex Ratio = 2.37). The standardized mortality rate for nasopharyngeal cancer in the world was 0.7 per 100,000 (1.0 per 100,000 in men and 0.4 per 100,000 in women)¹¹. The five countries that had the highest number of deaths from nasopharyngeal cancer included China with 21,300 deaths¹², Indonesia with 7,391 deaths, Vietnam with 2,885 deaths, India with 2,836 deaths, and Thailand with 1114 cases, respectively¹³. One of the main causes of death in Indonesia is nasopharyngeal cancer and at the time of initial diagnosis, 80% of the patients are at an advanced stage of the disease¹⁶. In Singapore, older patients with a diagnosis at stage 2 or stage 3 are at higher risk of recurrence and lower overall survival⁹. Nasopharyngeal cancers

TABLE 1. Factors related to the nasopharyngeal cancer.

Modifiable and environmental risk factors	Genetic or unmodifiable risk factors
Epstein-Barr virus (EBV)	Family history of NPC
HLA class I genotypes	Age
Salt-preserved fish	Sex
Lack of fresh fruits and vegetables	Socioeconomic status
Chronic respiratory tract conditions	
Tobacco smoke	
Herbal medicines	
Occupational dusts	
Formaldehyde	

are treated by radiation, but the result of treatment varies widely throughout the world. 8% of new cases have occurred in countries with poor treatment outcome. Global inequality in access to and the need for optimal services planning shows a meaningful relationship between survival and access to radiation therapy⁶. Over the past two decades, the treatment of nasopharyngeal cancer has improved significantly with the simultaneous introduction of chemotherapy and radiotherapy. The overall incidence of patients with metastases has remained about 25-34% and the survival of these patients is low^{21,22}. Lack of knowledge of general practitioners working in Asian medical centers may lead to a delay in diagnosis of nasopharyngeal cancer¹⁶.

RISK FACTORS FOR NASOPHARYNGEAL CANCER

Table 1 shows factors related to the nasopharyngeal cancer (based on modifiable or unmodifiable risk factors). The most important risk factors for nasopharyngeal cancer are presented in Table 2. Then, based on studies, the risk factors have been discussed.

SUMMARY OF POSSIBLE RISK FACTORS FOR NASOPHARYNGEAL CANCER

EPSTEIN-BARR VIRUS (EBV)

In many studies, a very strong association has been shown between nasopharyngeal carcinoma and EBV infection²³⁻²⁶. Approximately about 30 versions of the EBV gene are in the nucleus of malignant cells. Most versions are “mini circular

chromosomes”, called episomes. These episomes are in some cases adjacent to viral DNA versions²⁷. This association leads to the use of new methods using viral serological tests to make diagnosis and screening in the population at risk²⁸. High levels of IgA antibodies with the capsid antigen of EBV and early antigen (EA) in areas with high incidence of nasopharynx are a valuable screening tool²⁴.

INHERITANCE AND ETHNICITY

People with a family history of cancer, in particular a family history of nasopharyngeal cancer, are at risk more than 4 to 10 times for nasopharyngeal cancer^{4,29,30}. Studies⁴ conducted in Southern China revealed the role of a genetic combination with environmental effects in the development of nasopharyngeal cancer. In Southern China, there are three main ethnic groups: Cantonese, Hokkien-Teochiu (Minan-Chaoshan) and Hakka. The highest incidence of nasopharyngeal cancer was seen among Cantonese and the lowest among Hokkien-Teochiu and Hakka³¹. The remarkable feature of ethnic distribution in the Cantonese population is the genetic factor, but the combined role of genetic with the particular lifestyle and environmental factors should not be ignored³².

SALT-PRESERVED FISH AND OTHER FOODS

Many studies³²⁻³⁴ have documented the consumption of salt-preserved fish as one of the risk factors for the cancer. In Chinese populations, the relative risk of nasopharyngeal carcinoma in weekly consumers is lower than those who use salt-preserved fish very little or at all. Overall, it was about 1.4 to 3.2. Relative risk varied from 1.8 to 7.5 for those who consu-

TABLE 2. Summary of possible risk factors for NPC.

Factor	Strength of association	Consistency of association	Subgroup-specific associations
EBV	Strong	Consistent	More consistent association with types II and III NPC
Family history of NPC	Strong	Consistent	
HLA class I genotypes	Moderate to strong	Consistent	Inconsistent associations with HLA class II genotypes
Salt-preserved fish	Moderate to strong	Consistent	Stronger association with consumption at weaning
Lack of fresh fruits and vegetables	Moderate	Fairly consistent	
Chronic respiratory tract conditions	Moderate	Fairly consistent	
Tobacco smoke	Weak to moderate	Fairly consistent	Stronger association with type I NPC
Herbal medicines	Weak to moderate	Inconsistent	
Occupational dusts	Weak to moderate	Inconsistent	More consistent association with wood dust exposure
Formaldehyde	Weak to moderate	Inconsistent	



med daily³⁵⁻⁴⁰. The risk of nasopharyngeal cancer in relation to other preservative foods, including meat, eggs, fruits and vegetables, increases in Southern China, Southeast Asia, North Africa/Middle East, and Arctic natives^{37,41-43}. These preservative foods are also involved in people in areas with a low incidence in Northern China and the United States^{44,45}.

HUMAN LEUKOCYTE ANTIGEN GENES

Gene-related research on nasopharyngeal cancer focuses on human leukocyte antigen genes⁴⁶. In people with a weak HLA allele in the presence of EBV, the antigens likely increase the risk of nasopharyngeal cancer. In people with a strong HLA allele in the presence of EBV, the risk for this cancer is lower^{47,48}. In the meta-analysis on the Southern population of China, there was the evidence of a positive association of nasopharyngeal cancer with increased risk in HLA-A2, B14, B46 and a reverse association with HLA-A11, B13 and B22⁴⁹.

FRESH VEGETABLES AND FRUITS

Sufficient and adequate consumption of fresh fruits and vegetables is associated with a 30-50% reduction in the risk of nasopharyngeal cancer³². In contrast, lack of enough vegetables and fruits increases the risk of developing nasopharyngeal cancer⁵⁰. The results of a meta-analysis study showed that high consumption of vegetables is associated with a 36% reduction in the risk of nasopharyngeal cancer⁴³.

HISTORY OF RESPIRATORY DISEASES

Most studies^{37,40,51-58} showed that the risk of developing nasopharyngeal cancer in people with chronic rhinitis, sinusitis, nasal polyps, or ear infection, is almost twice. These findings indicate that benign inflammation and respiratory infection may make nasopharynx mucus susceptible to this cancer. In addition, some bacteria can reduce the conversion of nitrate to nitrite, which can then form N-nitroso carcinogenic compounds⁵⁹.

ALCOHOL CONSUMPTION

Alcohol consumption is associated with nasopharyngeal cancer in a complicated way. Many studies³² have shown that there is no strong evidence for the association between alcohol consumption and the risk of nasopharyngeal cancer. The results of a meta-analysis recently showed that high alcohol consumption is associated with a significant increase in the risk of nasopharyngeal cancer⁶⁰.

CIGARETTE SMOKING

Many studies^{30,42,51,61-67} have highlighted that cigarette smoking is associated with nasopharyngeal cancers. The pattern of the relationship between smoking and the risk of nasopharyngeal cancer is

dose-dependent^{51,63}. The results of a meta-analysis showed that the risk of the cancer in smokers was 60% higher than non-smokers⁶⁸.

OCCUPATIONAL EXPOSURE

Exposure to wood dust is known to be a risk factor for nasopharyngeal cancer^{64,69-72}. The results of a study on 29,000 wood workers in the UK and the United States showed the association between the exposure to wood dust and the increased risk of nasopharyngeal carcinoma⁷³. Several studies⁷⁴ have shown that exposure to formaldehyde increases the risk of nasopharyngeal cancer. In addition, IARC investigations in 2005⁷⁵ and two meta-analyses^{76,77} refer to the dose-response relationship between formaldehyde and the risk of developing nasopharyngeal cancer. Exposure to other chemicals or stimulants, such as steam, smoke and chemicals, flammable products, cotton dust⁵² or solvents such as phenoxy acid and chlorophenol, causes an increased risk of nasopharyngeal cancer^{51,78}.

HERBAL REMEDIES

Several studies have reported the association between the use of herbal medicines and the increased risk of nasopharyngeal cancer. Hildesheim et al⁷⁹ in a case-control study conducted in the Philippines, asked to subjects about ever use of any herbal medicines, and noted a statistically significant 2.5-fold risk among ever users. However, the latter study is difficult to interpret for the following reasons: recall bias is a serious concern when a nonspecific question (such as general use of herbal medicines without naming specific formulations) was asked under a case-control setting. In addition, the use of herbal medicines is part of the 'traditional' lifestyle, an established risk factor for NPC in Chinese as well as Southeast Asians. In other words, the use of herbal medicine may simply be a marker of the NPC-related lifestyle⁷⁹⁻⁸³. The consumption of slow-cooked (several hours) and canton style herbal teas are associated with reduced risk of nasopharyngeal cancer⁸⁴.

AGE AND SEX

The incidence of nasopharyngeal cancer in men is 2 to 3 times higher than women^{66,85}. Differences between men and women may be due to different lifestyle habits (e.g., tobacco consumption) or biological differences³². In most low-risk groups, the incidence of nasopharyngeal cancer is consistent with increasing age⁸⁶⁻⁸⁸. In contrast, in high-risk groups, the incidence increases in age group of 50 to 59 years, and then decreases⁸⁹. This is due to the fact that these groups are exposed to carcinogenic agents in the early stages of life⁹⁰. Nasopharyngeal cancer may take several decades to develop malignant cells, and then signs appear. Therefore, exposure to carcinogens in early life may have significant effects on the incidence of this cancer³².

SOCIOECONOMIC STATUS

Among the populations at risk for nasopharyngeal disease, including South China, natives of Southeast Asia and the Arabs of North Africa, the lower social class is associated with a higher risk of nasopharyngeal cancer^{41,91,92}. The consumption of stored food is known to be one of the most important risk factors for nasopharyngeal cancer. Given that these foods are cheap, consumption of these foods is higher for people with low socioeconomic status. Relationship between the social class and the risk factors for nasopharyngeal cancer, such as exposure to dust and smoking in the United States, have been reported⁽⁶⁶⁾.

CONCLUSIONS

We aimed at determining the incidence and mortality rate of nasopharyngeal cancer in the world and the relationship between environmental risk factors and the incidence of nasopharyngeal cancer. The findings of this study showed that the standardized incidence of nasopharyngeal cancer in the world was 1.2 per 100,000, and the standardized mortality rate for nasopharyngeal carcinoma was 0.7 per 100,000. The disease is more common in Southeast Asia. The most important risk factors for nasopharyngeal cancer include: EBV, heredity, human leukocyte antigen (HLA), salt-preserved fish, and history of respiratory diseases. Fruit and vegetable consumption is known as a protective agent for this cancer. Training programs, early screening, good life promotion policies, timely treatment and diagnosis are good for reducing the burden of this cancer.

CONFLICT OF INTEREST:

The authors declared no competing interests.

REFERENCES

1. TSAO SW LK, HUANG DP. NASOPHARYNGEAL CARCINOMA. In: TSELIS AC, JENSON H, EDITORS. Epstein-Bar virus. New York: Taylor & Francis. 2006; pp. 273-295.
2. RAAB-TRAU N. Epstein-Bar virus in the pathogenesis of NPC. In: Robertson ES, editor. Epstein-Barr virus Wymondham. Norfolk: Caister Academic Press 2005; pp. 71-92.
3. BUSSON P, KERYSER C, OOKA T, CORBEX M. EBV-associated nasopharyngeal carcinomas: from epidemiology to virus-targeting strategies. Trends Microbiol 2004; 12: 356-360.
4. CHANG ET, ADAMI HO. The enigmatic epidemiology of nasopharyngeal carcinoma. Cancer Epidemiol Biomarkers Prev 2006; 15: 1765-1777.
5. FERLAY J, SHIN H, BRAY F, FORMAN D, MATHERS C, PARKIN D. Incidence/mortality data. GLOBOCAN 2008 v2.0. Cancer incidence and mortality worldwide: IARC CancerBase No. 10 [Internet]. Lyon, France: International Agency for Research on Cancer, 2010.
6. LAM KO, LEE AW, CHOI CW, SZE HC, ZIETMAN AL, HOPKINS KI, ROSENBLATT E. Global pattern of nasopharyngeal cancer: correlation of outcome with access to radiation therapy. Int J Radiat Oncol Biol Phys 2016; 94: 1106-1112.
7. WEE J, HA TC, LOONG S, QIAN CN. Is nasopharyngeal cancer really a "Cantonese cancer"? Chin J Cancer 2010; 29: 517-526.
8. CHUA ML, WEE JT, HUI EP, CHAN AT. Nasopharyngeal carcinoma. Lancet 2016; 387: 1012-1024.
9. MAK HW, LEE SH, CHEE J, THAM I, GOH BC, CHAO SS, ONG YK, LOH KS, LIM CM. Clinical outcome among nasopharyngeal cancer patients in a multi-ethnic society in Singapore. PLoS One 2015; 10: e0126108.
10. XU ZJ, ZHENG RS, ZHANG SW, ZOU XN, CHEN WQ. Nasopharyngeal carcinoma incidence and mortality in China in 2009. Chin J Cancer 2013; 32: 453-458.
11. TANG LL, CHEN WQ, XUE WQ, HE YQ, ZHENG RS, ZENG YX, JIA WH. Global trends in incidence and mortality of nasopharyngeal carcinoma. Cancer Lett 2016; 374: 22-30.
12. CHEN W, ZHENG R, ZHANG S, ZENG H, XIA C, ZUO T, YANG Z, HE J. Cancer incidence and mortality in China, 2013. Cancer Lett 2017; 10: 63-71.
13. MAHDAVIFAR N, TOWHIDI F, MAKHSOSI BR, PAKZAD R, MOINI A, AHMADI A, LOTFI S, SALEHINIYA H. Incidence and mortality of nasopharynx cancer and its relationship with human development index in the world in 2012. World J Oncol 2016; 7: 109-118.
14. TEO MC, SOO KC. Cancer trends and incidences in Singapore. Jpn J Clin Oncol 2013; 43: 219-224.
15. VALEAN S, ACALOVSCI M, DICULESCU M, MANUC M, GOLDIS A, SFARTI C, TRIFAN A. Mortality in Digestive Cancers, 2012: International Data and Data from Romania. J Gastrointest Liver Dis 2015; 24: 507-514.
16. FLES R, WILDEMAN MA, SULISTIONO B, HARYANA SM, TAN IB. Knowledge of general practitioners about nasopharyngeal cancer at the Puskesmas in Yogyakarta, Indonesia. BMC Med Educ 2010; 10: 81-87.
17. CURADO M-P, EDWARDS B, SHIN HR, STORM H, FERLAY J, HEANUE M, BOYLE P. Cancer incidence in five continents, Volume IX: IARC Press, International Agency for Research on Cancer, 2007.
18. JEMAL A, BRAY F, CENTER MM, FERLAY J, WARD E, FORMAN D. Global cancer statistics. CA Cancer J Clin 2011; 61: 69-90.
19. ADHAM M, STOKER S, WILDEMAN M, RACHMADI L, GONDHOWIARDJO S, ATMAKUSUMAH D, GATOT D, FLES R, GREJER A, HERMANI B, MIDDELDORP J, TAN I. Current status of cancer care for young patients with nasopharyngeal carcinoma in Jakarta, Indonesia. PLoS One 2014; 9: e102353.
20. WILDEMAN MA, FLES R, HERDINI C, INDRASARI RS, VINCENT AD, TJOKRONAGORO M, STOKER S, KURNIANDA J, KARAKULLUKCU B, TAROENO-HARIADI KW, HAMMING-VRIEZE O, MIDDELDORP JM, HARIWIYANTO B, HARYANA SM, TAN IB. Primary treatment results of nasopharyngeal carcinoma (NPC) in Yogyakarta, Indonesia. PLoS One 2013; 8:e63706.
21. CHEN MY, JIANG R, GUO L, ZOU X, LIU Q, SUN R, QIU F, XIA ZJ, HUANG HQ, ZHANG L, HONG MH, MAI HQ, QIAN CN. Locoregional radiotherapy in patients with distant metastases of nasopharyngeal carcinoma at diagnosis. Chin J Cancer 2013; 32: 604-613.
22. LEE AW, LIN JC, NG WT. Current management of nasopharyngeal cancer. Semin Radiat Oncol 2012; 22: 233-244.
23. RICKINSON AB, KIEFF E. Epstein-Barr virus. In: Knipe DM, Howley PM, editors. Field's virology. 4th ed. Philadelphia (Pennsylvania): Lippincott, Williams & Wilkins; 2001; pp. 2575-2627.
24. LICITRA L, BERNIER J, CVITKOVIC E, GRANDI C, SPINAZZÉ S, BRUZZI P, GATTA G, MOLINARI R. Cancer of the nasopharynx. Crit Rev Oncol Hematol 2003; 45: 199-214.



25. LUNG ML, CHEUNG AKL, KO JMY, LUNG HL, CHENG Y, DAI W. The interplay of host genetic factors and Epstein-Barr virus in the development of nasopharyngeal carcinoma. *Chin J Cancer* 2014; 33: 556-562.
26. RAZAK AR, SIU LL, LIU F-F, ITO E, O'SULLIVAN B, CHAN K. Nasopharyngeal carcinoma: the next challenges. *Eur J Cancer* 2010; 46: 1967-1978.
27. KRIPALANI-JOSHI S, LAW H. Identification of integrated Epstein-Barr virus in nasopharyngeal carcinoma using pulse field gel electrophoresis. *Int J Cancer* 1994; 56: 187-192.
28. COHEN E, LEE A. Editorial for special issue on nasopharynx cancer. *Oral Oncol* 2014; 50: 325-330.
29. CHEN DL, HUANG TB. A case-control study of risk factors of nasopharyngeal carcinoma. *Cancer Lett* 1997; 117: 17-22.
30. EK BURANAWAT W, EK PANYASKUL C, BRENNAN P, KANKA C, TEPSUWAN K, TEMIYASTITH S, KLINVIMOL T, PONGNIKORN S, SANGRAJRANG S. Evaluation of non-viral risk factors for nasopharyngeal carcinoma in Thailand: results from a case-control study. *Asian Pac J Cancer Prev* 2010; 11: 929-932.
31. ARMSTRONG R, KUTTY MK, ARMSTRONG M. Self-specific environments associated with nasopharyngeal carcinoma in Selangor, Malaysia. *Soc Sci Med* 1978; 12: 149-156.
32. JIA WH, QIN HD. Non-viral environmental risk factors for nasopharyngeal carcinoma: a systematic review. *Semin Cancer Biol* 2012; 22: 117-126.
33. PUTERA I, RAMADHAN MG, ANINDYA S, SUTANTO NR, KURNIAWAN A, HOSEA FN, SAFITRI ED. Relationship between salted fish consumption and nasopharyngeal carcinoma: an evidence-based case report. *Acta Med Indones* 2015; 47: 72-77.
34. ADHAM M, KURNIAWAN AN, MUHTADI AI, ROEZIN A, HERMANI B, GONDHOWIARDJO S, TAN IB, MIDDELDORP JM. Nasopharyngeal carcinoma in Indonesia: epidemiology, incidence, signs, and symptoms at presentation. *Chin J Cancer* 2012; 31: 185-196.
35. HENDERSON B, LOUIE E. Discussion of risk factors for nasopharyngeal carcinoma. *IARC Sci Publ* 1977; 20: 251-260.
36. MIMI CY, HO JH, LAI SH, HENDERSON BE. Cantonese-style salted fish as a cause of nasopharyngeal carcinoma: report of a case-control study in Hong Kong. *Cancer Res* 1986; 46: 956-961.
37. YU MC, HUANG TB, HENDERSON BE. Diet and nasopharyngeal carcinoma: a case-control study in Guangzhou, China. *Int J Cancer* 1989; 43: 1077-1082.
38. ARMSTRONG RW, IMREY PB, LYE MS, ARMSTRONG MJ, YU MC, SANI S. Nasopharyngeal carcinoma in Malaysian Chinese: salted fish and other dietary exposures. *Int J Cancer* 1998; 77: 228-235.
39. YUAN JM, WANG XL, XIANG YB, GAO YT, ROSS RK, YU MC. Preserved foods in relation to risk of nasopharyngeal carcinoma in Shanghai, China. *Int J Cancer* 2000; 85: 358-363.
40. ZOU J, SUN Q, AKIBA S, YUAN Y, ZHA Y, TAO Z, WEI L, SUGAHARA T. A case-control study of nasopharyngeal carcinoma in the high background radiation areas of Yangjiang, China. *J Radiat Res* 2000; 41: 53-62.
41. SRIAMPORN S, VATANASAPT V, PISANI P, YONGCHAIYUDHA S, RUNGPIRARANGSRI V. Environmental risk factors for nasopharyngeal carcinoma: a case-control study in northeastern Thailand. *Cancer Epidemiol Biomarkers Prev* 1992; 1: 345-348.
42. CHENG YJ, HILDESHEIM A, HSU MM, CHEN IH, BRINTON LA, LEVINE PH, CHEN CJ, YANG CS. Cigarette smoking, alcohol consumption and risk of nasopharyngeal carcinoma in Taiwan. *Cancer Causes Control* 1999; 10: 201-207.
43. GALLICCHIO L, MATANOSKI G, TAO XG, CHEN L, LAM TK, BOYD K, ROBINSON KA, BALICK L, MICKELSON S, CAULFIELD LE, HERMAN JG, GUALLAR E, ALBERG AJ. Adulthood consumption of preserved and nonpreserved vegetables and the risk of nasopharyngeal carcinoma: a systematic review. *Int J Cancer* 2006; 119: 1125-1135.
44. NING JP, YU MC, WANG QS, HENDERSON BE. Consumption of salted fish and other risk factors for nasopharyngeal carcinoma (NPC) in Tianjin, a low-risk region for NPC in the People's Republic of China. *J Natl Cancer Inst* 1990; 82: 291-296.
45. FARROW DC, VAUGHAN TL, BERWICK M, LYNCH CF, SWANSON GM, LYON JL. Diet and nasopharyngeal cancer in a low-risk population. *Int J Cancer* 1998; 78: 675-679.
46. LI X, FASANO R, WANG E, YAO KT, MARINCOLA FM. HLA associations with nasopharyngeal carcinoma. *Curr Mol Med* 2009; 9: 751-765.
47. SIMONS M, WEE G, CHAN S, SHANMUGARATNAM K, DAY N. Immunogenetic aspects of nasopharyngeal carcinoma (NPC) III. HL-A type as a genetic marker of NPC predisposition to test the hypothesis that Epstein-Barr virus is an etiological factor in NPC. *IARC Sci Publ* 1975; 11: 249-258.
48. HILDESHEIM A, APPLE RJ, CHEN CJ, WANG SS, CHENG YJ, KLITZ W, MACK SJ, CHEN IH, HSU MM, YANG CS, BRINTON LA, LEVINE PH, ERLICH HA. Association of HLA class I and II alleles and extended haplotypes with nasopharyngeal carcinoma in Taiwan. *J Natl Cancer Inst* 2002; 94: 1780-1789.
49. GOLDSMITH D, WEST T, MORTON R. HLA associations with nasopharyngeal carcinoma in Southern Chinese: a meta-analysis. *Clin Otolaryngol Allied Sci* 2002; 27: 61-67.
50. KEY TJ, ALLEN NE, SPENCER EA, TRAVIS RC. The effect of diet on risk of cancer. *Lancet* 2002; 360: 861-868.
51. YU MC, GARABRANT DH, HUANG TB, HENDERSON BE. Occupational and other non-dietary risk factors for nasopharyngeal carcinoma in Guangzhou, China. *Int J Cancer* 1990; 45: 1033-1039.
52. HENDERSON BE, LOUIE E, JING JS, BUELL P, GARDNER MB. Risk factors associated with nasopharyngeal carcinoma. *N Engl J Med* 1976; 295: 1101-1106.
53. GESER A, CHARNAY N, DAY N, HO H. Environmental factors in the etiology of nasopharyngeal carcinoma: report on a case-control study in Hong Kong. *IARC Sci Publ* 1978; 20: 213-229.
54. CLIFFORD P. External Influences on the nose and throat: carcinogens in the nose and throat: nasopharyngeal carcinoma in Kenya. *Proc Roy Soc Med* 1972; 65: 682-686.
55. LIN TM, CHEN KP, LIN CC, HSU MM, TU SM, CHIANG TC, JUNG PF, HIRAYAMA T. Retrospective study on nasopharyngeal carcinoma 2. *J Natl Cancer Inst* 1973; 51: 1403-1408.
56. MIRABELLI MC, HOPPIN JA, TOLBERT PE, HERRICK RF, GNEPP DR, BRANN EA. Occupational exposure to chlorophenol and the risk of nasal and nasopharyngeal cancers among US men aged 30 to 60. *Am J Ind Med* 2000; 37: 532-541.
57. ZHU K, LEVINE RS, BRANN EA, HALL HI, CAPLAN LS, GNEPP DR. Case-control study evaluating the homogeneity and heterogeneity of risk factors between sinonasal and nasopharyngeal cancers. *Int J Cancer* 2002; 99: 119-123.

58. SHANMUGARATNAM K, TYE C, GOH E, CHIA K. Etiological factors in nasopharyngeal carcinoma: a hospital-based, retrospective, case-control, questionnaire study. *IARC Sci Publ* 1978; 20: 199-212.
59. BARTSCH H, OHSHIMA H, PIGNATELLI B, CALMELS S. Endogenously formed N-nitroso compounds and nitrosating agents in human cancer etiology. *Pharmacogenetics* 1992; 2: 272-277.
60. MARRON M, BOFFETTA P, ZHANG ZF, ZARIDZE D, WÜNSCH-FILHO V, WINN DM, WEI Q, TALAMINI R, SZESZENIA-DABROWSKA N, STURGIS EM, SMITH E, SCHWARTZ SM, RUDNAI P, PURDUE MP, OLSHAN AF, ELUF-NETO J, MUSCAT J, MORGENSTERN H, MENEZES A, MCCLEAN M, MATOS E, MATES IN, LISSOWSKA J, LEVI F, LAZARUS P, LA VECCHIA C, KOIFMAN S, KELSEY K, HERRERO R, HAYES RB, FRANCESCHI S, FERNANDEZ L, FABIANOVA E, DAUDT AW, DAL MASO L, CURADO MP, CADONI G, CHEN C, CASTELLSAGUE X, BOCCIA S, BENHAMOU S, FERRO G, BERTHILLER J, BRENNAN P, MOLLER H, HASHIBE M. Cessation of alcohol drinking, tobacco smoking and the reversal of head and neck cancer risk. *Int J Epidemiol* 2009; 39: 182-196.
61. YUAN JM, WANG XL, XIANG YB, GAO YT, ROSS RK, YU MC. Non-dietary risk factors for nasopharyngeal carcinoma in Shanghai, China. *Int J Cancer* 2000; 85: 364-369.
62. JI X, ZHANG W, XIE C, WANG B, ZHANG G, ZHOU F. Nasopharyngeal carcinoma risk by histologic type in central China: impact of smoking, alcohol and family history. *Int J Cancer* 2011; 129: 724-732.
63. VAUGHAN TL, SHAPIRO JA, BURT RD, SWANSON GM, BERWICK M, LYNCH CF, LYON JL. Nasopharyngeal cancer in a low-risk population: defining risk factors by histological type. *Cancer Epidemiol Biomarkers Prev* 1996; 5: 587-593.
64. ARMSTRONG RW, IMREY PB, LYE MS, ARMSTRONG MJ, YU MC, SANI S. Nasopharyngeal carcinoma in Malaysian Chinese: occupational exposures to particles, formaldehyde and heat. *Int J Epidemiol* 2000; 29: 991-998.
65. NEŠIĆ V, ŠIPETIĆ S, VLAJINAC H, STOŠIĆ-DIVJAK S, JEŠIĆ S. Risk factors for the occurrence of undifferentiated carcinoma of nasopharyngeal type: a case-control study. *Srp Arh Celok Lek* 2010; 138: 6-10.
66. YU MC, YUAN JM. Epidemiology of nasopharyngeal carcinoma. *Semin Cancer Biol* 2002; 12: 421-429.
67. LUBIN JH, PURDUE M, KELSEY K, ZHANG ZF, WINN D, WEI Q, TALAMINI R, SZESZENIA-DABROWSKA N, STURGIS EM, SMITH E, SHANGINA O, SCHWARTZ SM, RUDNAI P, NETO JE, MUSCAT J, MORGENSTERN H, MENEZES A, MATOS E, MATES IN, LISSOWSKA J, LEVI F, LAZARUS P, LA VECCHIA C, KOIFMAN S, HERRERO R, FRANCESCHI S, WÜNSCH-FILHO V, FERNANDEZ L, FABIANOVA E, DAUDT AW, MASO LD, CURADO MP, CHEN C, CASTELLSAGUE X, BRENNAN P, BOFFETTA P, HASHIBE M, HAYES RB. Total exposure and exposure rate effects for alcohol and smoking and risk of head and neck cancer: a pooled analysis of case-control studies. *Am J Epidemiol* 2009; 170: 937-947.
68. XUE WQ, QIN HD, RUAN HL, SHUGART YY, JIA WH. Quantitative association of tobacco smoking with the risk of nasopharyngeal carcinoma: a comprehensive meta-analysis of studies conducted between 1979 and 2011. *Am J Epidemiol* 2013; 178: 325-338.
69. ARMSTRONG RW, ARMSTRONG MJ, MIMI CY, HENDERSON BE. Salted fish and inhalants as risk factors for nasopharyngeal carcinoma in Malaysian Chinese. *Cancer Res* 1983; 43: 2967-2970.
70. YANG XR, DIEHL S, PFEIFFER R, CHEN CJ, HSU WL, DOSEMECI M, CHENG YJ, SUN B, GOLDSTEIN AM, HILDESHEIM A; Chinese and American Genetic Epidemiology of NPC Study Team. Evaluation of risk factors for nasopharyngeal carcinoma in high-risk nasopharyngeal carcinoma families in Taiwan. *Cancer Epidemiol Biomarkers Prev* 2005; 14: 900-905.
71. VAUGHAN TL, STEWART PA, TESCHKE K, LYNCH CF, SWANSON GM, LYON JL, BERWICK M. Occupational exposure to formaldehyde and wood dust and nasopharyngeal carcinoma. *Occup Environ Med* 2000; 57: 376-384.
72. HILDESHEIM A, DOSEMECI M, CHAN CC, CHEN CJ, CHENG YJ, HSU MM, CHEN IH, MITTL BF, SUN B, LEVINE PH, CHEN JY, BRINTON LA, YANG CS. Occupational exposure to wood, formaldehyde, and solvents and risk of nasopharyngeal carcinoma. *Cancer Epidemiol Biomarkers Prev* 2001; 10: 1145-1153.
73. DEMERS PA, BOFFETTA P, KOGEVINAS M, BLAIR A, MILLER BA, ROBINSON CF, ROSCOE RJ, WINTER PD, COLIN D, MATOS E, VAINIO H. Pooled reanalysis of cancer mortality among five cohorts of workers in wood-related industries. *Scand J Work Environ Health* 1995; 21: 179-190.
74. PUÑAL-RIOBÓO J, VARELA-LEMA L, BARROS-DIOS JM, JUIZ-CRESPO MA, RUANO-RAVIÑA A. La ocupación como factor de riesgo del cáncer oral y de faringe. *Acta Otorrinolaringol Esp* 2010; 61: 375-383.
75. COGLIANO VJ, GROSSE Y, BAAN RA, STRAIF K, SECRETAN MB, EL GHISASSI F; Working Group for Volume 88. Meeting report: summary of IARC monographs on formaldehyde, 2-butoxyethanol, and 1-tert-butoxy-2-propanol. *Environ Health Perspect* 2005; 113: 1205-1208.
76. BLAIR A, SARACCI R, STEWART PA, HAYES RB, SHY C. Epidemiologic evidence on the relationship between formaldehyde exposure and cancer. *Scand J Work Environ Health* 1990; 16: 381-393.
77. PARTANEN T. Formaldehyde exposure and respiratory cancer – a meta-analysis of the epidemiologic evidence. *Scand J Work Environ Health* 1993; 19: 8-15.
78. HARDELL L, JOHANSSON B, AXELSON O. Epidemiological study of nasal and nasopharyngeal cancer and their relation to phenoxy acid or chlorophenol exposure. *Am J Ind Med* 1982; 3: 247-257.
79. HILDESHEIM A, WEST S, DEVEYRA E, DE GUZMAN MF, JURADO A, JONES C, IMAI J, HINUMA Y. Herbal medicine use, Epstein-Barr virus, and risk of nasopharyngeal carcinoma. *Cancer Res* 1992; 52: 3048-3051.
80. WEST S, HILDESHEIM A, DOSEMECI M. Non-viral risk factors for nasopharyngeal carcinoma in the philippines: results from a case-control study. *Int J Cancer* 1993; 55: 722-727.
81. LIN T, YANG C, TU S, CHEN C, KUO K, HIRAYAMA T. Interaction of factors associated with cancer of the nasopharynx. *Cancer* 1979; 44: 1419-1423.
82. ZHENG Y, TUPPIN P, HUBERT A, JEANNEL D, PAN YJ, ZENG Y, DE THÉ G. Environmental and dietary risk factors for nasopharyngeal carcinoma: a case-control study in Zangwu County, Guangxi, China. *Br J Cancer* 1994; 69: 508-514.
83. CHELLENG P, NARAIN K, DAS H, CHETIA M, MAHANTA J. Risk factors for cancer nasopharynx: a case-control study from Nagaland, India. *Natl Med J India* 2000; 13: 6-8.
84. JIA W-H, LUO XY, FENG BJ, RUAN HL, BEI JX, LIU WS, QIN HD, FENG QS, CHEN LZ, YAO SY, ZENG YX. Traditional Cantonese diet and nasopharyngeal carcinoma risk: a large-scale case-control study in Guangdong, China. *BMC Cancer* 2010; 10: 446-451.



85. **PARKIN D, WHELAN SL, FERLAY J, TEPPLO L, THOMAS DB.** Cancer incidence in five continents. IARC Scientific Publications No. 155. France 2002; pp. 773.
86. **WANG H, SEOW A, LEE HP.** Trends in cancer incidence among Singapore Malays: a low-risk population. *Ann Acad Med Singapore* 2004; 33: 57-62.
87. **LUO J, CHIA KS, CHIA SE, REILLY M, TAN CS, YE W.** Secular trends of nasopharyngeal carcinoma incidence in Singapore, Hong Kong and Los Angeles Chinese populations, 1973-1997. *Eur J Epidemiol* 2007; 22: 513-521.
88. **SUN LM, EPPLEIN M, LI CI, VAUGHAN TL, WEISS NS.** Trends in the incidence rates of nasopharyngeal carcinoma among Chinese Americans living in Los Angeles County and the San Francisco metropolitan area, 1992-2002. *Am J Epidemiol* 2005; 162: 1174-1178.
89. **LEE AW, FOO W, MANG O, SZE WM, CHAPPELL R, LAU WH, KO WM.** Changing epidemiology of nasopharyngeal carcinoma in Hong Kong over a 20-year period (1980-99): an encouraging reduction in both incidence and mortality. *Int J Cancer* 2003; 103: 680-685.
90. **SHANMUGARATNAM K. NASOPHARYNX.** IN: **SCHOTTENFELD D, FRAUMENI JF, JR.,** EDITORS. *Cancer epidemiology and prevention.* Philadelphia: W. B. Saunders Company 1982; pp. 536-553.
91. **MIMI CY, HO JH, ROSS RK, HENDERSON BE.** Nasopharyngeal carcinoma in Chinese – salted fish or inhaled smoke? *Prev Med* 1981; 10: 15-24.
92. **JEANNEL D, HUBERT A, DE VATHAIRE F, ELLOUZ R, CAMOUN M, BEN SALEM M, SANCHO-GARNIER H, DE-THÉ G.** Diet, living conditions and nasopharyngeal carcinoma in Tunisia – a case-control study. *Int J Cancer* 1990; 46: 421-425.