



LETTER TO THE EDITOR

ASSOCIATION BETWEEN STATIN USE AND INCIDENCE OF BREAST CANCER

A. SAFARI¹, H. SALEHINIYA², K. ALLAH BAKESHEI³, A. MOHAMMADIAN-HAFSHEJANI^{4,5}

¹Gonabad University of Medical Sciences, Gonabad, Iran

²Zabol University of Medical Sciences, Zabol, Iran

³Department of Social Medicine, School of Public Health, Dezfoul University of Medical Sciences, Dezfoul, Iran

⁴Department of Epidemiology and Biostatistics, School of Public Health, Shahrekord University of Medical Sciences, Shahrekord, Iran

⁵Department of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

KEYWORDS: *Statin, Incidence, Breast Cancer.*

LIST OF ABBREVIATIONS: : *BC: Breast cancer, CI: Confidence Interval.*

Dear Editor,

Breast cancer (BC) is the most common cancer in women. A total of 252,710 new cases and 40610 deaths from BC occurred in women in the United States in 2017. The risk of BC was 12.5% in the American women's lifetime¹. The pathway and risk factors for this cancer have not been fully diagnosed. Obesity and overweight, family history of BC, genetic factors, environmental factors, low physical activity, and exposure to chemical are among the diagnosed BC risk factors^{2,3}. A lot of studies have been planned to identify substances with a chemopreventive potential against breast cancer. The pharmaceutical group of Statins is widely prescribed in the world with the aim to prevent cardiovascular events and stroke.

The protective effects of Statins have been proven against the onset of some cancers. However, controversial results⁴ have been observed in studies on the effects of Statins on the incidence of BC, so that some separate investigations have considered Statins as the protective agents^{5,6}, as

risk factors⁷⁻⁹ and neutral¹⁰. There is not any definitive conclusion on the effects of Statins on the risk of BC. Undoubtedly, the results of meta-analysis studies can be helpful in this regard. Several meta-analysis researches on this field have not found any significant relationship between Statin use and the increase or decrease in the BC risk, so that the relative risk of BC in the treatment group compared to the control group was obtained equal to 0.94 (CI of 95%, 0.86-1.03) in the research by Islam et al¹¹, equal to 1.02 (CI: 95%, 0.95-1.09) in research by Qi-Jun et al¹², equal to 0.99 (CI of 95%, 0.94-1.04) in study by Undela et al⁴, and equal to 1.02 (CI of 95%, 0.89-1.18) in study by Bonova et al¹⁰. According to a research by Islam et al¹¹, the relative risk of BC was equal to 0.94 (CI of 95%, 0.70-1.28) for those who received Statins for 5 years or less and equal to 0.74 (CI of 95%, 0.52-1.04) for those who received Statins for over 5 years¹¹.

Furthermore, there was not any significant difference between Statins and the risk of BC in different parts of the world, so that the relative risk of BC in the treatment group compared to the control group was equal to 0.85 (CI of 95%, 0.41-1.52) in Asia, 1.01 (CI of 95%, 0.97-1.05) in Europe, 1.13 (CI of 95%, 0.44-2.92) in Oceania, and 0.92 (CI of 95%, 0.84-1.02) in the United States¹¹. Therefore, based on the results of meta-analysis studies, which



provide the highest level of scientific inferences, the use of statin does not significantly increase or decrease the risk of BC in recipients. Furthermore, the incidence of BC does not depend on the duration of Statin use and the geographical area of residence. Consequently, statins can be used in prevention of cardiovascular diseases and stroke without any worry about the increased risk of BC.

CONFLICT OF INTERESTS:

The Authors declare that they have no conflict of interests.

REFERENCES

1. SIEGEL RL, MILLER KD, JEMAL A. Cancer Statistics, 2017. *CA Cancer J Clin* 2017; 67: 7-30.
2. MAJEED W, ASLAM B, JAVED I, KHALIQ T, MUHAMMAD F, ALI A, RAZA A. Breast cancer: major risk factors and recent developments in treatment. *Asian Pac J Cancer Prev* 2014; 15: 3353-3358.
3. ROJAS K, STUCKEY A. Breast cancer epidemiology and risk factors. *Clin Obstet Gynecol* 2016; 59: 651-672.
4. UNDELA K, SRIKANTH V, BANSAL D. Statin use and risk of breast cancer: a meta-analysis of observational studies. *Breast Cancer Res Treat* 2012; 135: 261-269.
5. POCOBELLI G, NEWCOMB PA, TRENTAM-DIETZ A, TITUS-ERNSTOFF L, HAMPTON JM, EGAN KM. Statin use and risk of breast cancer. *Cancer* 2008; 112: 27-33.
6. HAUKKA J, SANKILA R, KLAUKKA T, LONNOVIST J, NISKANEN L, TANSKANEN A, WAHLBECK K, TIHONEN J. Incidence of cancer and statin usage--record linkage study. *Int J Cancer* 2010; 126: 279-284.
7. BECK P, WYSOWSKI DK, DOWNEY W, BUTLER-JONES D. Statin use and the risk of breast cancer. *J Clin Epidemiol* 2003; 56: 280-285.
8. EATON M, EKLOF J, BEAL JR, SAHMOUN AE. Statins and breast cancer in postmenopausal women without hormone therapy. *Anticancer Res* 2009; 29: 5143-5148.
9. COOGAN PF, ROSENBERG L, PALMER JR, STROM BL, ZAUBER AG, SHAPIRO S. Statin use and the risk of breast and prostate cancer. *Epidemiology* 2002; 13: 262-267.
10. BONOVAS S, FILIOUSI K, TSAVARIS N, SITARAS NM. Use of statins and breast cancer: a meta-analysis of seven randomized clinical trials and nine observational studies. *J Clin Oncol* 2005; 23: 8606-8612.
11. ISLAM MM, YANG HC, NGUYEN PA, POLY TN, HUANG CW, KEKADA S, KHALFAN AMB, DEBNATH T, LI YJ, ABDUL SS. Exploring association between statin use and breast cancer risk: an updated meta-analysis. *Arch Gynecol Obstet* 2017; 296: 1043-1053.
12. WU QJ, TU C, LI YY, ZHU J, QIAN KQ, LI WJ, WU L. Statin use and breast cancer survival and risk: a systematic review and meta-analysis. *Oncotarget* 2015; 6: 42988-43004.